arXiv:cond-mat/0202237v1 [cond-mat.mes-hall] 14 Feb 2002

Hyperfine-mediated transitions between a Zeeman split doublet in GaAs quantum dots: The role of the internal field

Sigurdur I. Erlingsson, Yuli V. Nazarov
Delft University of Technology, Department of Applied Physics, Lorentzweg 1, 2628 Delft, The Netherlands
(Dated: February 1, 2008)

We consider the hyperfine-mediated transition rate between Zeeman split spin states of the lowest orbital level in a GaAs quantum dot. We separate the hyperfine Hamiltonian into a part which is diagonal in the orbital states and another one which mixes different orbitals. The diagonal part gives rise to an effective (internal) magnetic field which, in addition to an external magnetic field, determines the Zeeman splitting. Spin-flip transitions in the dots are induced by the orbital mixing part accompanied by an emission of a phonon. We evaluate the rate for different regimes of applied magnetic field and temperature. The rates we find are bigger that the spin-orbit related rates provided the external magnetic field is sufficiently low.

PACS numbers: 71.23.La, 71.70.Jp

I. INTRODUCTION

The electron spin states in bulk semiconductor and heterostructures have attracted much attention in recent years. Experiments indicate very long spin decoherence times and small transition rates between states of different spin[1,2,3]. These promising results have motivated proposals for information processing based on electron spins in quantum dots, which might lead to a realization of a quantum computer[4,5].

A quantum dot is region where electrons are confined. The energy spectrum is discrete, due to the small size, and can display atomic-like properties[6,7]. Here we will consider quantum dots in GaAs-AlGaAs heterostructures. The main reasons for studying them are that relevant quantum dots are fabricated in such structures and Ga and As nuclei have a substantial hyperfine interactions with the conduction electrons. There are two main types of gate controlled dots in these systems, so-called vertical and lateral dots[8]. They differ in transverse confinement, which is approximately a triangular well and a square well for the lateral and vertical dots, respectively.

Manipulation the electron spin while maintaining phase coherence requires that it should be relatively well isolated from the environment. Coupling a quantum dot, or any closed quantum system, to its environment can cause decoherence and dissipation. One of the measures of the strength of the coupling to the environment are the transition rates, or inverse lifetimes, between the quantum dot states. In GaAs there are two main mechanisms that can cause finite lifetimes of spin states. These are the spin-orbit interaction and the hyperfine interaction with the surrounding nuclei. If a magnetic field is applied the change in Zeeman energy accompanying a spin-flip has to be absorbed by phonon emission. For two electron quantum dots, where the transitions are between triplet and singlet spin states, both spin-orbit[9,10] and hyperfine[11] mediated transitions have been studied. In both cases the transition rates are much smaller than the usual phonon rates, the spin-orbit rate being the higher except when the excited singlet and triplet states cross.[11] For low magnetic fields, i.e. away from the single-triplet transition, the energy of the emitted phonon can be quite large and the transitions involve deformation phonons rather than piezoelectric ones.

If there is odd number of electrons in the dot the ground-state is usually a spin doublet so that energy change associated with the spin-flip is the electron Zeeman energy. Owing to the small g-factor in GaAs this energy is rather small compared to the level spacing and the dominating phonon mechanism is due to piezoelectric phonons. Recently spin-orbit mediated spin-flip transitions between Zeeman levels was investigated[12]. Due to Kramer's degeneracy the transition amplitude for the spin-flip is proportional to the Zeeman splitting. This results in a spin-orbit spin-flip rate proportional to the fifth power of the Zeeman splitting.

In this publication we consider hyperfine mediated transitions between Zeeman split levels in a quantum dot. The transition amplitude remains finite at zero external magnetic field, giving a spin-flip rate (Eq. (27)) that is proportional to the cube of the Zeeman splitting. The cause of this is an internal magnetic field due to the hyperfine interaction. We consider the important concept of internal magnetic field in some detail. Since the parameters of hyperfine interaction between conduction band electrons and underlying nuclei in GaAs have been extensively investigated[13,14], including the Overhauser effect and spin-relaxation in GaAs/AlGaAs heterostructures[15,16,17,18,19] , we are able now to predict the typical time scale for this process in particular quantum dot geometries.

Upon completion of this work we learned about recent results of Khaetskii, Loss and Glazman[20]. They consider essentially the same situation and model and obtain *electron spin decoherence* without considering any mechanism of dissipation. This is in clear distinction from the present result for *spin-flip rate* that requires a mechanism of dissipation, i.e. phonons.

The rest of the paper is organized as follows: in section II the model used is introduced in addition to the basic assumptions and approximations used, section III

deals with the internal magnetic field due to the hyperfine interaction and section IV contains the derivation of the transition rates. Finally, in section V the results are discussed.

II. MODEL AND ASSUMPTIONS

We consider a quantum dot embedded in a AlGaAs/GaAs heterostructure. The details of the quantum dot eigenstates are not important for now, it suffices to say that the energy spectrum is discrete and the wave functions are localized in space. The spatial extension of the wavefunction in the lateral and transverse direction (growth direction) are denoted with ℓ and z_0, respectively. Quantum dots in these heterostructures are formed at a GaAs/AlGaAs interface, where the confining potential is very strong so that $\ell \gg z_0$. We define the volume occupied by an electron as $V_{\rm QD} = \pi\ell^2 z_0$. The Hamiltonian of the quantum dot can be written on the form

$$H_0 = \sum_l \big(\varepsilon_l + g\mu_B \boldsymbol{B}\cdot\hat{\boldsymbol{S}}\big)|l\rangle\langle l|, \tag{1}$$

where ε_l are the eigenenergies which depend on the structure of the confining potential and the applied magnetic field. The magnetic field also couples to the electron spin via the Zeeman term.

Since the Γ point of the conduction band in GaAs is mainly composed of s orbitals the dipole interaction with the nuclei vanishes and the hyperfine interaction can be described by the usual contact term

$$H_{\rm HF} = A\hat{\boldsymbol{S}}\cdot\sum_k \hat{\boldsymbol{I}}_k \delta(\boldsymbol{r}-\boldsymbol{R}_k), \tag{2}$$

where $\hat{\boldsymbol{S}}$ ($\hat{\boldsymbol{I}}_k$) and $\boldsymbol{r}$ ($\boldsymbol{R}_k$) denotes, respectively, the spin and position of the electron (kth nuclei). The delta function indicates the point-like nature of the contact interaction will result in a position dependant coupling. The coupling constant A has the dimension Volume×Energy. To get a notion of the related energy scale, it is straightforward to relate A to the energy splitting of the doublet for a fully polarized nuclear system,

$$E_n = AC_nI, \tag{3}$$

C_n being density of nuclei and I is the spin of a nucleus. In GaAs this energy is $E_n \approx 0.135\,\text{meV}$, which corresponds to a magnetic field of about $5\,\text{T}$.[15] For a given quantum dot geometry the number of nuclei occupying the dot is defined as $N_{\rm QD} = C_nV_{\rm QD}$. Since the Ga and As nuclei have the same spin and their coupling constants are comparable, we will assume that all the nuclear sites are characterized by the same hyperfine coupling A, and the $C_n \approx a_0^{-3}$ where a_0 is the lattice spacing. For realistic quantum dots $N_{\rm QD} \approx 10^4 - 10^6$, and it is therefore an important big parameter in the problem.

The coupling between the electron and the phonon bath is represented with

$$H_{\rm ph} = \sum_{\boldsymbol{q},\nu} \alpha_\nu(\boldsymbol{q})(b_{\boldsymbol{q},\nu}e^{i\boldsymbol{q}\cdot\boldsymbol{r}} + b^\dagger_{\boldsymbol{q},\nu}e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}), \tag{4}$$

where $b^\dagger_{\boldsymbol{q}\nu}$ and $b_{\boldsymbol{q}\nu}$ are creation and annihilation operator for the phonon mode with wave-vector $\boldsymbol{q}$ on brach ν. In GaAs there are two different coupling mechanisms, deformation ones and piezoelectric ones. For transitions between Zeeman split levels in GaAs, i.e. low energy emission, the most effective phonon mechanism is due to piezoelectric phonons. We will assume that the heterostructure is grown in the [100] direction. This is the case for almost all dots and it imposes important symmetry relations on the coupling coefficient. The square of the coupling coefficient for the piezoelectric phonons is then given by (see Ref. 21)

$$\alpha^2_\nu(\boldsymbol{q}) = \frac{(eh_{14})^2\hbar}{2\rho c_\nu V q} A_\nu(\theta) \tag{5}$$

where (eh_{14}) is the piezoelectric coefficient, ρ is the mass density, c_ν is the speed of sound of branch ν, V is normalization volume and we have defined $\boldsymbol{q} = q(\cos\phi\sin\theta, \sin\phi\sin\theta, \cos\theta)$ and the A_ν's are the so-called anisotropy functions, see appendix B.

III. INTERNAL MAGNETIC FIELD

In this section we will introduce the concept of the effective magnetic field, the *internal field*, acting on the electron due to the hyperfine interaction. This internal field is a semiclassical approximation to the nuclear system, this approximation being valid in the limit of large number of nuclei, $N_{\rm QD} \gg 1$. If the nuclei are noticeably polarized, this field almost coincides with the Overhauser field that represents the average nuclear polarization. It is important that the internal field persists even at zero polarization giving rise to doublet splitting of the order $E_nN_{\rm QD}^{-1/2}$.

First we write the Hamiltonian in Eq. (2) in the basis of the electron orbital states and present it as a sum of two terms

$$H_{\rm HF} = H^0_{\rm HF} + V_{\rm HF}, \tag{6}$$

where the terms are defined as

$$H^0_{\rm HF} = A\sum_l |l\rangle\langle l|\hat{\boldsymbol{S}}\cdot\sum_k |\langle \boldsymbol{R}_k|l\rangle|^2\hat{\boldsymbol{I}}_k \tag{7}$$

$$V_{\rm HF} = \sum_{l\neq l'} |l\rangle\langle l'|\hat{\boldsymbol{S}}\cdot\sum_k \langle l|\boldsymbol{R}_k\rangle\langle\boldsymbol{R}_k|l'\rangle\hat{\boldsymbol{I}}_k. \tag{8}$$

By definition $H^0_{\rm HF}$ does not couple different orbital levels. By combining Eqs. (1) and (7) one obtains the following Hamiltonian

$$H_0 = \sum_l \big(\varepsilon_l + g\mu_B\hat{\boldsymbol{\mathcal{B}}}_l\cdot\hat{\boldsymbol{S}}\big)|l\rangle\langle l|. \tag{9}$$

We will regard the mixing term $V_{\rm HF}$ as a perturbation to H_0. The justification for this is that the typical fluctuations of the electron energy due to the hyperfine interaction are much smaller than the orbital energy separation, $\hbar\Omega$.

We now concentrate on H_0 and formulate semiclassical description of it. For this we consider the operator of the orbitally dependent effective magnetic field;

$$\hat{\boldsymbol{\mathcal{B}}}_l = \boldsymbol{B} + \frac{1}{g\mu_B}\hat{\boldsymbol{K}}_l \tag{10}$$

where

$$\hat{\boldsymbol{K}}_l = \frac{E_n}{IC_n}\sum_k |\langle \boldsymbol{R}_k|l\rangle|^2 \hat{\boldsymbol{I}}_k. \tag{11}$$

Our goal is to replace the operator $\hat{\boldsymbol{K}}_l$ with a classical field. To prove the replacement is reasonable we calculate the average of the square for a given unpolarized nuclear state $|\boldsymbol{\mu}\rangle$

$$\boldsymbol{K}_l^2 = \langle \boldsymbol{\mu}|\hat{\boldsymbol{K}}_l^2|\boldsymbol{\mu}\rangle = \frac{E_n^2}{N_{\rm QD}}\rho_l^{[2]}(I+1)/I \tag{12}$$

where $\rho_l^{[n]} = \int d\boldsymbol{r} V_{\rm QD}^{n-1}|\langle \boldsymbol{r}|l\rangle|^{2n}$ is a dimensionless constant depending on the wave function of the orbital l. We cannot simply replace $\hat{\boldsymbol{K}}_l$ by its eigenvalues: as in the case of the usual spin algebra, different components of $\hat{\boldsymbol{K}}_l$ do not commute. In addition its square does not commute with individual components, $[\hat{\boldsymbol{K}}_l^2, K_l^\alpha] \neq 0$. To estimate fluctuations of $\hat{\boldsymbol{K}}_l$ we calculate the uncertainty relations between its components

$$\Delta K_l^\alpha \Delta K_l^\beta \geq \frac{E_n^2}{N_{\rm QD}^{3/2}} = \boldsymbol{K}_l^2 \frac{1}{N_{\rm QD}^{1/2}}. \tag{13}$$

Since $N_{\rm QD}^{-1/2} \ll 1$ we have proved that the quantum fluctuations in $\hat{\boldsymbol{K}}_l$ are much smaller than its typical amplitude. The semiclassical picture introduced above is only valid for high temperatures, $kT \gg E_n N_{\rm QD}^{-1}$, where there are many states available to the nuclear system and the typical length is proportional to $N_{\rm QD}^{1/2}$. For temperatures below $E_n N_{\rm QD}^{-1}$ the nuclear system will predominantly be in the ground state and the classical picture breaks down. This is similar to the quantum mechanical description of a particle moving in a potential. At zero temperature it will be localized in some potential minimum and quantum mechanics will dominate. At sufficiently high temperatures the particle occupies higher energy states and its motion is well described by classical mechanics. Having established this we can replace the average over the density matrix of the nuclei by the average over a classical field $\boldsymbol{K}_l$, note that it has no 'hat', whose values are Gaussian distributed

$$P(\boldsymbol{K}_l) = \left(\frac{3}{2\pi\langle \boldsymbol{K}_l^2\rangle}\right)^{3/2} \exp\Big(-\frac{3(\boldsymbol{K}_l - \boldsymbol{K}_l^{(0)})^2}{2\langle \boldsymbol{K}_l^2\rangle}\Big). \tag{14}$$

Here $\boldsymbol{K}_l^{(0)}$ is the average, or Overhauser, field assuming small degree of polarization. The effective magnetic field acting on the electron is $\boldsymbol{\mathcal{B}_l} = \mathcal{B}_l \boldsymbol{n}$ where $\boldsymbol{n}$ is the unit vector along the total field for a given $\boldsymbol{K}_l$, see Fig. 1. For this configuration the spin eigenfunctions are $|\boldsymbol{n}_\pm\rangle$ corresponding to eigenvalues

$$\boldsymbol{n}\cdot\hat{\boldsymbol{S}}|\boldsymbol{n}_\pm\rangle = \pm\frac{1}{2}|\boldsymbol{n}_\pm\rangle \tag{15}$$

The effective Zeeman Hamiltonian is then

$$H_{\rm Z} = g\mu_B \boldsymbol{\mathcal{B}}_l \cdot \hat{\boldsymbol{S}} \tag{16}$$

in this given internal field configuration. The spectrum of H_0 thus consists of many doublets distinguished by the value of $\boldsymbol{\mathcal{B}}_l$. The magnitude of the effective field $\mathcal{B}_l$ determines the Zeeman splitting of the each doublet

$$\begin{aligned}\Delta_l &= g\mu_B \mathcal{B}_l \\ &= (E_B^2 + K_l^2 + 2E_B K_l \cos\theta)^{1/2}\end{aligned} \tag{17}$$

where $E_B = g\mu_B B$ the external field Zeeman energy.

We conclude this section with two remarks concerning time and energy scales. First, since the dynamics of $\boldsymbol{K}_l$ is essentially the dynamics of the nuclear system it changes on a time scale of the nuclear relaxation, which can be quite long.[16] For electron dynamics on shorter timescales $\boldsymbol{K}_l$ plays the role of a 'frozen disorder'. At longer time scales self-averaging over all configurations of $\boldsymbol{K}_l$ takes place. Second, the RMS value of $\boldsymbol{K}_l$ is approximately $5\,\mu$eV which corresponds to a 100 Gauss field (for a value $N_{\rm QD} \approx 10^5$).

IV. TRANSITION RATE

We concentrate on the transitions between the doublet components in Eq. (15). Assuming that the higher energy doublet state is initially occupied, we will calculate the transition rate to the lower one. The transition must be accompanied by energy dissipation equal to Δ_0. This energy cannot be absorbed be the nuclear system.[11,22] So we need an external mechanism of energy dissipation. The most effective one is known to be phonons. However the phonons alone cannot change the electron spin so we need a mechanism which mixes spin and orbital degrees of freedom, that is $V_{\rm HF}$ from Eq. (8). Thus the transition amplitude is proportional to both $V_{\rm HF}$ and the electron phonon coupling $H_{\rm ph}$.

Here we assume that the electron is in the lowest orbital state $|0\rangle$ since the phonon mechanism will bring the electron to this state from any higher orbital on timescales much smaller than those related to transtions between the doublet components. Thus we consider an initial state of the entire system $|{\rm i}\rangle = |0, \boldsymbol{n}_-; \boldsymbol{\mu}; \boldsymbol{N}\rangle$ which is a product state of the electron, nuclear $|\boldsymbol{\mu}\rangle$ and phonon $|\boldsymbol{N}\rangle$ systems and the final state $|{\rm f}\rangle = |0, \boldsymbol{n}_+; \boldsymbol{\mu}'; \boldsymbol{N}'\rangle$. The

transition amplitude between them, in second order perturbation theory, reads

$$\mathcal{T} = \sum_{l\neq 0}\Bigg(\frac{\langle 0,\boldsymbol{n}_+;\boldsymbol{\mu}'|V_{\rm HF}|l,\boldsymbol{n}_-;\boldsymbol{\mu}\rangle\langle l;\boldsymbol{N}'|H_{\rm ph}|0;\boldsymbol{N}\rangle}{(\varepsilon_0-\varepsilon_l)+E_B} + \frac{\langle 0;\boldsymbol{N}'|H_{\rm ph}|l;\boldsymbol{N}\rangle\langle l,\boldsymbol{n}_+;\boldsymbol{\mu}'|V_{\rm HF}|0,\boldsymbol{n}_-;\boldsymbol{\mu}\rangle}{(\varepsilon_0-\varepsilon_l)-E_B}\Bigg) \quad (18)$$

The summation is over virtual states involving higher orbitals and the denominators in Eq. (18) contains the energy differences between different orbital states. The internal field depends on the orbital state, resulting in a rather complicated expression. Albeit the energy related to the internal field is much smaller than the orbital separation so we can safely replace the Zeeman splitting with E_B. The reason for that is that only at high external fields where $\Delta_0 \approx E_B$ will the effects of the Zeeman splitting be appreciable in the denominator. The internal field also appears in the phonon rate since it determines the electron energy difference between the initial and final state. Since $H_{\rm ph}$ does not connect different nuclear states and conversely $V_{\rm HF}$ does not mix different phonon states the sums over intermediate phonon and nuclear states reduce to a single term. From this transition amplitude the transition rate is obtained via Fermis golden rule

$$\tilde{\Gamma}_{\rm sf} = \frac{2\pi}{\hbar}\sum_{\boldsymbol{N}'}\sum_{\boldsymbol{\mu}'}|\mathcal{T}|^2\delta(E_{\rm i}-E_{\rm f}), \quad (19)$$

where $E_{\rm i}-E_{\rm f}$ is the energy difference between the initial and final state of the combined systems. Substituting Eq. (18) into Eq. (19) we get the following relation for the spin-flip rate

$$\begin{aligned}\tilde{\Gamma}_{\rm sf} = \sum_{l,l'\neq 0}\Bigg(&\frac{\langle l,\boldsymbol{n}_-;\boldsymbol{\mu}|V_{\rm HF}|0,\boldsymbol{n}_+\rangle\langle 0,\boldsymbol{n}_+|V_{\rm HF}|l',\boldsymbol{n}_-;\boldsymbol{\mu}\rangle}{((\varepsilon_0-\varepsilon_l)+E_B)((\varepsilon_0-\varepsilon_{l'})+E_B)}\frac{2\pi}{\hbar}\sum_{\boldsymbol{N}'}\langle 0;\boldsymbol{N}|H_{\rm ph}|l;\boldsymbol{N}'\rangle\langle l';\boldsymbol{N}'|H_{\rm ph}|0;\boldsymbol{N}\rangle\delta(E_{\rm i}-E_{\rm f})\\
&+\frac{\langle l,\boldsymbol{n}_-;\boldsymbol{\mu}|V_{\rm HF}|0,\boldsymbol{n}_+\rangle\langle l',\boldsymbol{n}_+|V_{\rm HF}|0,\boldsymbol{n}_-;\boldsymbol{\mu}\rangle}{((\varepsilon_0-\varepsilon_l)+E_B)((\varepsilon_0-\varepsilon_{l'})-E_B)}\frac{2\pi}{\hbar}\sum_{\boldsymbol{N}'}\langle 0;\boldsymbol{N}|H_{\rm ph}|l;\boldsymbol{N}'\rangle\langle 0;\boldsymbol{N}'|H_{\rm ph}|l';\boldsymbol{N}\rangle\delta(E_{\rm i}-E_{\rm f})\\
&+\frac{\langle 0,\boldsymbol{n}_-;\boldsymbol{\mu}|V_{\rm HF}|l,\boldsymbol{n}_+\rangle\langle 0,\boldsymbol{n}_+|V_{\rm HF}|l',\boldsymbol{n}_-;\boldsymbol{\mu}\rangle}{((\varepsilon_0-\varepsilon_l)-E_B)((\varepsilon_0-\varepsilon_{l'})+E_B)}\frac{2\pi}{\hbar}\sum_{\boldsymbol{N}'}\langle l;\boldsymbol{N}|H_{\rm ph}|0;\boldsymbol{N}'\rangle\langle l';\boldsymbol{N}'|H_{\rm ph}|0;\boldsymbol{N}\rangle\delta(E_{\rm i}-E_{\rm f})\\
&+\frac{\langle 0,\boldsymbol{n}_-;\boldsymbol{\mu}|V_{\rm HF}|l,\boldsymbol{n}_+\rangle\langle l',\boldsymbol{n}_+|V_{\rm HF}|0,\boldsymbol{n}_-;\boldsymbol{\mu}\rangle}{((\varepsilon_0-\varepsilon_l)-E_B)((\varepsilon_0-\varepsilon_{l'})-E_B)}\frac{2\pi}{\hbar}\sum_{\boldsymbol{N}'}\langle l;\boldsymbol{N}|H_{\rm ph}|0;\boldsymbol{N}'\rangle\langle 0;\boldsymbol{N}'|H_{\rm ph}|l';\boldsymbol{N}\rangle\delta(E_{\rm i}-E_{\rm f})\Bigg). \quad (20)\end{aligned}$$

The spin-flip rate depends on the initial state of the nuclear system $|\boldsymbol{\mu}\rangle$. This poses the problem of how to deal with the nuclear state $|\boldsymbol{\mu}\rangle$, since we already demoted all the spin operators to a collective classical variable. A conceptually simple solution lies in the fact that when Eq. (20) and (8) are considered together one sees that the rate is a sum over all pairs of nuclei in the system. Focusing on a given pair of nuclei k and k', all the other nuclei are unchanged when the electron spin is 'scattered' on by this pair. By simply redefining the classical field such that it is composed of all nuclei except this given pair we can circumvent the problem. This procedure will not change our previous result regarding the properties of $\boldsymbol{K}_l$ and also defining $|\boldsymbol{\mu}\rangle = |m_k\rangle|m_{k'}\rangle$ makes it straightforward to work with the nuclear states in Eq. (20).

Although the transition rate can be very slow, the typical duration of a transition event is set by energy uncertainty, $\hbar/\Delta_0$. This is much shorter than the typical time for nuclear spin relaxation so that the nuclear system is frozen in a given configuration of $\boldsymbol{K}_0$. In this case taking an average over $\boldsymbol{K}_0$ is not well motivated since a given configuration of $\boldsymbol{K}_0$ is fixed. For now we will postpone the average over the classical field. After performing a thermal average over $|\boldsymbol{\mu}\rangle$ in the transition rate we obtain the following

$$\begin{aligned}\Gamma_{\rm sf} = G_{\rm corr}\Bigg(&\sum_{l\neq 0}\Bigg\{\frac{2a_{ll}\gamma_{ll}}{\delta\varepsilon_l^2}\left(1+3\frac{E_B^2}{\delta\varepsilon_l^2}\right)+\frac{2\Re\{\tilde{a}_{ll}\tilde{\gamma}_{ll}\}}{\delta\varepsilon_l^2}\left(1+\frac{E_B^2}{\delta\varepsilon_l^2}\right)\Bigg\}\\
+\sum_{l<l'\neq 0}\Bigg\{&\frac{4\Re\{a_{ll'}\gamma_{ll'}\}}{\delta\varepsilon_l\delta\varepsilon_{l'}}\Big(1+\Big(\frac{(\delta\varepsilon_l+\delta\varepsilon_{l'})^2}{\delta\varepsilon_l\delta\varepsilon_{l'}}-1\Big)\frac{E_B^2}{\delta\varepsilon_l\delta\varepsilon_{l'}}\Big)+\frac{4\Re\{\tilde{a}_{ll'}\tilde{\gamma}_{ll'}\}}{\delta\varepsilon_l\delta\varepsilon_{l'}}\Big(1+\Big(\frac{(\delta\varepsilon_l-\delta\varepsilon_{l'})^2}{\delta\varepsilon_l\delta\varepsilon_{l'}}+1\Big)\frac{E_B^2}{\delta\varepsilon_l\delta\varepsilon_{l'}}\Big)\Bigg\}\Bigg) \quad (21)\end{aligned}$$

where $\delta\varepsilon_l = \varepsilon_0 - \varepsilon_l$. Here we have introduced the one-site nuclear spin correlation function $G_{\rm corr}$, which is evaluated in appendix A. The denominator has been expanded to second order in the Zeeman splitting. The parameters $a_{ll'}, \tilde{a}_{ll'}$ are related to the $V_{\rm HF}$ matrix elements

$$a_{ll'} = A^2 C_n \int d^3\boldsymbol{R}_k \Psi_l^*(\boldsymbol{R}_k)|\Psi_0(\boldsymbol{R}_k)|^2\Psi_{l'}(\boldsymbol{R}_k) \quad (22)$$

$$\tilde{a}_{ll'} = A^2 C_n \int d^3\boldsymbol{R}_k \Psi_l^*(\boldsymbol{R}_k)\Psi_{l'}^*(\boldsymbol{R}_k)\Psi_0(\boldsymbol{R}_k)^2 \quad (23)$$

and $\gamma_{ll'}, \tilde{\gamma}_{ll'}$ are generalized phonon transition rates

$$\gamma_{ll'} = \frac{2\pi}{\hbar}\sum_{\boldsymbol{q}\nu} \frac{\alpha_\nu^2(\boldsymbol{q})[e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}]_{0,l}[e^{i\boldsymbol{q}\cdot\boldsymbol{r}}]_{l',0}}{e^{\beta\hbar\omega_{\boldsymbol{q},\nu}}-1}\delta(\hbar\omega_{\boldsymbol{q}\nu}-\Delta_0) \quad (24)$$

$$\tilde{\gamma}_{ll'} = \frac{2\pi}{\hbar}\sum_{\boldsymbol{q}\nu} \frac{\alpha_\nu^2(\boldsymbol{q})[e^{-i\boldsymbol{q}\cdot\boldsymbol{r}}]_{0,l}[e^{i\boldsymbol{q}\cdot\boldsymbol{r}}]_{0,l'}}{e^{\beta\hbar\omega_{\boldsymbol{q},\nu}}-1}\delta(\hbar\omega_{\boldsymbol{q}\nu}-\Delta_0) \quad (25)$$

Here we have only included the emission process since we assume the the spin is initially in the higher energy state.

Until now we have considered a general quantum dot and the rate in Eq. (21) is valid for any quantum dot. To proceed further we will specify the confining potential to be parabolic in the lateral direction and in the transverse one a triangular well potential is chosen. The wave function is $\langle \boldsymbol{r}|l\rangle \equiv \chi_0(z)\psi_{n,M}(r,\theta)$ where n, M denote the orbital and angular momentum quantum numbers respectively of the Darwin-Fock solution and $\chi_0(z)$ is the wave function in the transverse direction. The square of the lateral confining length is $\ell^2 = \hbar^2/m^*\hbar\Omega$, where m^* is the electron effective mass and $\Omega = (\Omega_0^2 + (\omega_c/2)^2)^{1/2}$ is the effective confining frequency. The confining energy is $\hbar\Omega_0$ and $\omega_c = eB/m^*$ is the cyclotron frequency. What remains is to calculate the a's and γ's in Eqs. (22)-(25). The results of these calculations are presented appendix B.

In principle it is possible to obtain the rate for all parameter values but to make the discussion more transparent we will consider regimes of the applies magnetic field (i) $E_B \approx E_n N_{\rm QD}^{-1/2}$ and (ii) $E_B \gg E_n N_{\rm QD}^{-1/2}$. In regime (i) both $\Delta_0 \ll \hbar c_\nu \ell^{-1}$ and $\Delta_0 \ll kT$ (for experimentally relevant temperatures) and only the lowest order terms in $\Delta_0/\hbar c_\nu \ell^{-1}$ need to be considered. In GaAs $\hbar c_l \ell^{-1} = 3.3\ell^{-1}\,\mathrm{nm}\times\mathrm{meV}$ and $\hbar c_t \ell^{-1} = 2.0\ell^{-1}\,\mathrm{nm}\times\mathrm{meV}$ for the longitudinal and transverse branches respectively. In the other regime we assume that the applied field dominates and that the internal field may be ignored, but no additional assumption are made in this case. This will in general require some numerical work. The resulting hyperfine-mediated spin-flip rates are

$$\Gamma_{\rm HF} = 0.34\frac{G_{\rm corr}}{I^2}\frac{E_n^2}{N_{\rm QD}(\hbar\Omega)^2}\frac{(eh_{14}\ell)^2 kT}{8\pi\rho c^5\hbar^4}\left(E_B^2 + K_0^2 + 2E_B K_0\cos\theta\right) \quad \text{for } E_B \approx E_n N_{\rm QD}^{-1/2} \quad (26)$$

$$\Gamma_{\rm HF} = \frac{G_{\rm corr}}{I^2}\frac{E_n^2(n(E_B)+1)}{N_{\rm QD}(\hbar\Omega)^2}\frac{(eh_{14}\ell)^2 E_B^3}{8\pi\rho c^5\hbar^4}\left(C_0(E_B) + \left(\frac{E_B}{\hbar\Omega}\right)^2 C_2(E_B)\right) \quad \text{for } E_B \gg E_n N_{\rm QD}^{-1/2} \quad (27)$$

Note that the rates have different dependence on the emitted energy Δ_0. In regime (i) it is proportional to Δ_0^2 due to the Bose statistics, and in regime (ii) it is proportional to Δ_0^3 for $\Delta_0 \ll \hbar c_\nu \ell^{-1}$. In Eq. (27) we introduce the functions C_0 and C_2 which contain the details of the higher orbitals and the anisotropy integrals. For low fields $\Delta_0 \ll \hbar c_\nu \ell^{-1}$ these functions are constant. The value for the spin flip rate in Eq. (26), for some typical value of $K_0 \approx 10^{-2}\,\mathrm{T}$, is very low $\Gamma_{\rm HF} \approx 10^{-6}\,\mathrm{s}$. This results in a lifetime of days, which will be extremely difficult to measure. We have plotted the general spin-flip rate in Eq. (27) for different confining energies and temperatures in Figs. 2 and 3, for both in-plane and perpendicular applied magnetic field. In the case of the higher confining energy there is very small difference between the in-plane and perpendicular direction of the external magnetic field. For the lower confining energy there is a substantial difference between the two directions of magnetic field. In this case the approximation $\Delta_0 \ll \hbar c_\nu \ell^{-1}$ is no longer good and the the Δ_0^3 dependence of the rate is changed by the C-functions. The values of the rates are quite small, depending on the applied field, being in $\sim 1\,\mathrm{s}^{-1}$ for $T = 4\,\mathrm{K}$ at $B \approx 0.5\,\mathrm{T}$ for a confining frequency of $\hbar\Omega_0 = 2\,\mathrm{meV}$.

V. DISCUSSION

Generally speaking spin-flip rates require external dissipation. This is why at small Zeeman splittings they will contain small factor reflecting the vanishing phonon density of states. For a spin-orbit rate[12], the Kramer's degeneracy results in this small factor being proportional to E_B^5. Presence of nuclear spins violate the Kramer's theorem. So that the hyperfine rate discussed in the present paper is proportional to E_B^3 and will dominate at sufficiently low fields.

Comparing the hyperfine rate in Eq. (27) to spin-orbit related rates[12] $\Gamma_{\rm SO}$ we obtain that the hyperfine rate dominates for fields below $B \approx 0.3\,\mathrm{T}$ (assuming

$z_0 = 10\,\mathrm{nm}$ and $\hbar\Omega_0 = 4\,\mathrm{meV}$).

The role of the internal field produced by the nuclei is that the spin-flip rate does not vanish even for in the absence of external magnetic field. The show that the minimum rate is rather small, corresponding to a relaxation time of the order of days. We believe that the internal field will play an important role when the full dynamics of the electron spin in the presence of the nuclear system is considered.

Our model should also be applicable to other polar semiconductors which have non-zero nuclear spin, e.g. InAs where the g-factor is much larger.

VI. AKNOWLEDGEMENT

One author (SIE) would like to thank Daniela Pfannkuche, Alexander Chudnovskii and Daniel Loss for fruitful discussions. This work is a part of the research program of the "Stichting vor Fundementeel Onderzoek der Materie (FOM)"

APPENDIX A: AVERAGE OVER NUCLEAR SPIN PAIR

When the thermal average over a pair of free nuclei the following nuclear correlation function appear in Eq. (21)

$$G_{\mathrm{corr}} = (S^{\alpha}_{+-})^* S^{\beta}_{+-} \langle \delta I^{\alpha}_k \delta I^{\beta}_{k'} \rangle_T = (S^{\alpha}_{+-})^* S^{\beta}_{+-} \langle \delta \hat{I}^{\alpha} \delta \hat{I}^{\beta} \rangle_T \delta_{k,k'}, \tag{A1}$$

where $\delta \hat{I}^{\alpha} = \hat{I}^{\alpha} - \langle I^{\alpha} \rangle_T$ and the electron spin matrix elements being $S^{\alpha}_{+-} = \langle \boldsymbol{n}_+ | S^{\alpha} | \boldsymbol{n}_- \rangle$. The kronekker delta reflects that there are no correlations between two different nuclei and we have dropped the k subscript in $\langle \delta \hat{I}^{\alpha} \delta \hat{I}^{\beta} \rangle_T$ since the nuclei is assumed to be identical. By defining the symmetric correlator

$$g^{\alpha\beta} = \frac{1}{2} \langle \delta \hat{I}^{\alpha} \delta \hat{I}^{\beta} + \delta \hat{I}^{\beta} \delta \hat{I}^{\alpha} \rangle_T \tag{A2}$$

we get the following

$$G_{\mathrm{corr}} = (S^{\alpha}_{+-})^* S^{\beta}_{+-} (g^{\alpha\beta} + i/2 \epsilon^{\alpha\beta\gamma} \langle \hat{I}^{\gamma} \rangle_T). \tag{A3}$$

In an isotropic system, $\langle \hat{\boldsymbol{I}} \rangle_T = 0$, the value of the correlation function is $G_{\mathrm{corr}} = \frac{1}{2}\frac{1}{3} I(I+1)$.

APPENDIX B: MATRIX ELEMENTS FOR A PARABOLIC QUANTUM DOT

Using the Darwin-Fock solutions and the Fang-Howard variational solution for the triangular quantum well we obtain the following equation are Eqs. (22)-(25)

$$a_{ll'}\gamma_{ll'} = \delta_{M,M'} \frac{A^2 C n}{V_{\mathrm{QD}} \zeta} \frac{(e h_{14} \ell)^2 \Delta^3}{8\pi\rho c^5 \hbar^4} (n(\Delta)+1) \frac{\Gamma(n+n'+|M|+1) 2^{-3(n+n'+|M|)}}{n! n'! (n+|M|)! (n'+|M|)!} \times \left(\sum_{\nu} \frac{c^5}{c_\nu^5} \left(\frac{\Delta \ell}{\hbar c_\nu} \right)^{2(n+n'+|M|-1)} \int_0^{\pi} d(\cos\theta) A_\nu(\theta) \frac{(\sin\theta)^{2(n+n'+|M|)} \exp\left(-\frac{1}{2} \left(\frac{\Delta \sin\theta}{\hbar c_\nu \ell^{-1}} \right)^2 \right)}{\left(1 + \left(\frac{\Delta \cos\theta}{3\hbar c_\nu z_0^{-1}} \right)^2 \right)^3} \right) \tag{B1}$$

where $\zeta^{-1} = z_0 \int dz |\chi(z)|^4$ and $c^{-5} = c_l^{-5} + c_t^{-5}$ is the effective sound velocity of the phonons. The anisotropy functions are

$$A_t(\theta) = \frac{\sin^2\theta (8\cos^4\theta + \sin^4\theta)}{4} \tag{B2}$$

$$A_l(\theta) = \frac{9\cos^2\theta \sin^4\theta}{2}. \tag{B3}$$

The equation for $\tilde{a}_{ll'}\tilde{\gamma}_{ll'}$ is identical except for a different Kronekker delta function $\delta_{M,-M'}$. The integral needs in general to be evaluated numerically but when $\Delta \ll \hbar c_\nu \ell^{-1}$ the exponential term and the denominator become unity and the resulting integral is simple to calculate.

[1] J. M. Kikkawa and D. D. Awschalom, Phys. Rev. Lett. **80**, 4313 (1997).
[2] Y. Ohno *et al.*, Physica E **6**, 817 (2000).
[3] T. Fujisawa, Y. Tokura, and Y. Hirayama, Phys. Rev. B **63**, R81304 (2001).
[4] D. Loss and D. P. DiVincenzo, Phys. Rev. A **57**, 120

(1998).
[5] G. Burkard, D. Loss, and D. P. DiVincenzo, Phys. Rev. B **59**, 2070 (1999).
[6] R. C. Ashoori, Nature **379**, 413 (1996).
[7] S. Tarucha *et al.*, Phys. Rev. Lett. **77**, 3613 (1996).
[8] L. P. Kouwenhoven *et al.*, in *Mesoscopic Electron Transport*, *NATO Series*, edited by L. L. Sohn, L. P. Kouwenhoven, and G. Schoen (Kluwer, Dordrecht, 1997).
[9] A. V. Khaetskii and Y. V. Nazarov, Phys. Rev. B **61**, 12639 (2000).
[10] A. V. Khaetskii and Y. V. Nazarov, Physica E **6**, 470 (2000).
[11] S. I. Erlingsson, Y. V. Nazarov, and V. I. Fal'ko, Phys. Rev. B **64**, 195306 (2001).
[12] A. V. Khaetskii, Physica E **10**, 27 (2001).
[13] D. Paget, G. Lampel, and B. Sapoval, Phys. Rev. B **15**, 5780 (1977).
[14] D. Paget and V. L. Berkovits, in *Optical Orientation*, edited by F. Meier and B. P. Zakharchenya (North-Holland, Amsterdam, 1984).
[15] M. Dobers *et al.*, Phys. Rev. Lett. **61**, 1650 (1988).
[16] A. Berg, M. Dobers, R. R. Gerhardts, and K. v. Klitzing, Phys. Rev. Lett. **64**, 2563 (1990).
[17] I. D. Vagner, T. Maniv, and E. Ehrenfreund, Solid State Comm. **44**, 635 (1982).
[18] S. E. Barrett *et al.*, Phys. Rev. Lett. **74**, 5112 (1995).
[19] R. Tycko *et al.*, Science **268**, 1460 (1995).
[20] A. V. Khaetskii, D. Loss, and L. Glazman, cond-mat/0201303 (2002).
[21] P. J. Price, Ann. Phys. **133**, 217 (1981).
[22] J. H. Kim, I. D. Vagner, and L. Xing, Phys. Rev. B **49**, 16777 (1994).



FIG. 1: The internal field coordinate system is set by the external magnetic field $\boldsymbol{B}$, i.e. $\boldsymbol{e}_z \parallel \boldsymbol{B}$. The combination of the external and the internal field $\boldsymbol{K}_l$ results in an effective field $\mathcal{B}_l$.



FIG. 2: The hyperfine-mediated spin-flip rate for a quantum dot with $z_0 = 10\,\mathrm{nm}$ and $\hbar\Omega_0 = 5\,\mathrm{meV}$, plotted a function of external magnetic field for two different temperatures $T = 0.1\,\mathrm{K}$ and $4\,\mathrm{K}$.



FIG. 3: The hyperfine-mediated spin-flip rate for a quantum dot with $z_0 = 10\,\mathrm{nm}$ and $\hbar\Omega_0 = 5\,\mathrm{meV}$, plotted a function of external magnetic field for two different temperatures $T = 0.1\,\mathrm{K}$ and $4\,\mathrm{K}$.